

# CMIP5 – Performance and Climate Change Assessment of maximum and minimum temperatures in Europe

M.J. Carvalho, P. Melo-Gonçalves and A. Rocha

CESAM and Department of Physics, University of Aveiro

Correspond to mariajcarvalho@ua.pt



## Motivation

Climate Change, especially in the regime of **extreme events** is the focus of many studies. This is due to the fact that extreme events have a near-immediate effect on every-day life issues.

Under a changing climate, it is expected that these events will become more common.

Is this true for Europe? And if so, what are the geographical differences?

## Objective

I. Evaluate the performance of CMIP5 models in recent-past simulations for **Europe**.

II. Within Europe, define areas of coherent climate change patterns in temperature

III. Determine what is happening and will happen in the future to extreme events.

## Model Performance

| Model               | tasmax/<br>tasmin |
|---------------------|-------------------|
| ACCESS1-0           | 99.6              |
| CanESM              | 99.3              |
| CCSM4               | 99.7              |
| CNRM-CM5            | 99.7              |
| CSIRO Mk3-6-0       | 99.6              |
| EC-EARTH r1i1p1     | 99.8              |
| EC-EARTH r12i1p1    | 99.8              |
| HadGEM2-ES          | 99.6              |
| GFDL-ESM2M          | 99.4              |
| IPSL-CM5A-MR        | 99.4              |
| MIROC5              | 99.7              |
| MIROC-ESM-CHEM      | 99.3              |
| MPI-ESM-LR (r1i1p1) | 99.6              |
| MPI-ESM-LR (r2i1p1) | 99.6              |
| NorESM1-M           | 99.5              |

Taking the time series for the recent-past (1986 - 2005), the 2 sample **Kolmogorov-Smirnov** test was used to compare the modeled and observed data sets for maximum and minimum temperatures, at the 95% confidence level.

All models perform similarly. However, in order to reduce uncertainty associated to climate change projections an ensemble of the mentioned models was used

Table 1: List of models and % of overland points where the maximum (tasmax)/ minimum

tasmin) were used. All models show the same distribution at the 95% confidence level.

## Regionalization

RCP8.5 Daily Climat. – Historical Daily Climat.

Multi-feature K-Means Clustering Analysis

The multi-features are, not only the 2 variables, but also the 15 simulations.




Figure 1: European Regions of coherent seasonal climate change for minimum and maximum temperature for the 13-model ensemble.

## Data & Methodology

### OBSERVED DATASET:

- E-OBS V9.0 on a 0.25 degree regular grid
- Re-gridded to each of the models' grids

### MODELED DATASETS:

- Model List in Table 1.
- Each model has a different horizontal resolution;
- Ensemble mean is determined based on the re-gridded form of the models, with a 1.5° x 1.5° horizontal resolution.

### TIME PERIODS:

- Recent-past 1986 – 2005
- Long-term future 2081 – 2100

**RCP 8.5**

### INDEXES USED:

- **Maximum Temperature**
  - **Cold Days (TX10p):** Number of days when the tasmax < 10<sup>th</sup> percentile.
  - **Warm Days (TX90p):** Number of days when the tasmax > 10<sup>th</sup> percentile.
- **Minimum Temperature**
  - **Cold Nights (TN10p):** Number of days when the tasmin < 10<sup>th</sup> percentile.
  - **Warm Nights (TN90p):** Number of days when the tasmin > 90<sup>th</sup> percentile.



The percentile was determined using a 5-day window, based on the recent-past period.

## Results & Conclusions




Figure 2: Mean yearly series of the cold/ warm days and cold/ warm nights for the recent-past (1986-2005) on the left and long-term future (2081-2100) on the right for each of the European regions.

### Model Evaluation:

Most models show the same overall performance level. There is little difference between results from the same model with different initializations.

### Recent-Past:

- decrease in cold days and cold nights
- increase in warm days and warm nights

Higher maximum and minimum temperatures.

### Long-Term Future:

- Cold nights and days tend to disappear;
- Warm nights and days, as defined tend towards the total number of days per year;
- If the long-term-future percentile was used instead of the recent-past, there would still be an increase in warm nights/ days and decrease in cold nights/ days, but not reaching the mentioned magnitudes.

This work was supported by FEDER funds through the Programa Operacional Factores de Competitividade – COMPETE and by Portuguese national funds through FCT – Fundação para a Ciência e a Tecnologia, within the framework of the RESORT Project Reference PTDC/CTE-ATM/111508/2009 and CLIPE Project Reference PTDC/AAC-CLI/111733/2009.